

Oxidative stress and condition-dependent sexual signals: more than just seeing red

Michael Garratt and Robert C. Brooks

Proc. R. Soc. B published online 30 May 2012
doi: 10.1098/rsob.2012.0568

References

[This article cites 94 articles, 31 of which can be accessed free](#)

<http://rsob.royalsocietypublishing.org/content/early/2012/05/24/rsob.2012.0568.full.html#ref-list-1>

P<P

Published online 30 May 2012 in advance of the print journal.

Subject collections

Articles on similar topics can be found in the following collections

[behaviour](#) (797 articles)

[evolution](#) (1206 articles)

[physiology](#) (26 articles)

Email alerting service

Receive free email alerts when new articles cite this article - sign up in the box at the top right-hand corner of the article or click [here](#)

Advance online articles have been peer reviewed and accepted for publication but have not yet appeared in the paper journal (edited, typeset versions may be posted when available prior to final publication). Advance online articles are citable and establish publication priority; they are indexed by PubMed from initial publication. Citations to Advance online articles must include the digital object identifier (DOIs) and date of initial publication.

Review

Oxidative stress and condition-dependent sexual signals: more than just seeing red

Michael Garratt* and Robert C. Brooks

*Evolution and Ecology Research Centre, and School of Biological, Earth and Environmental Sciences,
The University of New South Wales, Sydney, New South Wales 2052, Australia*

The links between fitness, health, sexual signals and mate choice are complex and subject to ongoing study. In 1999, von Schantz *et al.* made the valuable suggestion that oxidative stress may be an important missing piece of this complex puzzle. Their suggestion has been enthusiastically tested, with over 300 studies citing their paper, but most effort has concerned carotenoid-based (and to a lesser extent melanin-based) visual signals, predominantly in birds and fishes. Today, we know a great deal more about oxidative stress and related physiology, in both a pathological and regulatory sense, than we did in 1999. We revisit von Schantz *et al.*'s predictions and, more importantly, highlight novel mechanisms that could link oxidative stress with a range of energetically demanding signals, greatly increasing the scope from visual signalling systems that are usually discussed and nearly always tested. In particular, we argue that differences between individuals in their ability to regulate physiology related to oxidative stress may be an important factor influencing the production of sexual signals and the costs that are incurred from investment.

Keywords: oxidative stress; sexual selection; metabolism; energy production; redox regulation

1. INTRODUCTION

Sexual signals involved in the attraction of mates show great variety, from the elaborate plumage and complex song of many birds to the odours, mating rituals and acoustic calls employed by mammals and invertebrates. Despite this diversity, a trend predominates; high-quality signallers frequently produce the largest, brightest or most elaborate signals and potential mates typically choose these signallers over those with smaller, duller or less elaborate signals. For such signalling systems to evolve and remain stable, signals must reliably and honestly indicate those signallers that provide direct or genetic benefits as mates [1]. For simplicity's sake, the vast majority of research in this area has concerned male 'signallers' and female 'choosers'.

Honest signalling can be maintained if high- and low-quality individuals differ in some kind of physiological state, usually referred to as 'condition', that relates to health, vigour, acquired resources and ultimately reproductive value. Condition-dependence either constrains the ability of low-quality individuals to produce and maintain sexual signals [2] or generates differential costs from signalling investment [3]. Evolutionary biologists have a tendency to treat the physiological factors linking sexual signals to condition as a 'black box' [4], or refer to condition purely in relation to resource acquisition and allocation [5].

Thirteen years ago, von Schantz *et al.* [6] suggested that one particular aspect of physiology—oxidative stress—may link 'the expression of sexual ornaments to genetic variation in fitness-related traits, thus promoting

the evolution of female mate choice and male sexual ornamentation'. More recently, evolutionary biologists have shown even greater interest in oxidative stress, suggesting that it could be a key physiological factor generating life-history trade-offs and constraining investment in a whole range of traits, such as growth, reproduction and lifespan [7–9]. The role of oxidative stress in life-history evolution has gained interest from researchers working on variety of different taxa. However, researchers working on oxidative stress and sexual selection have focused almost entirely on species that produce visual signals, such as birds and fishes, despite von Schantz *et al.*'s original prediction that oxidative stress could maintain honesty in a range of signalling systems. One reason for this narrow focus relates to the proposed mechanisms that could link oxidative stress with sexual signalling, as some of these pathways are applicable only to species that produce visual sexual signals. Some mechanisms have also been falsified over the past 13 years or shown to be based on an outdated understanding of oxidative stress. In this article, we briefly review these potential mechanisms in the light of modern knowledge of oxidative stress. More importantly, we use this knowledge to highlight other greatly underappreciated physiological pathways that may maintain the honesty of a range of sexual signals, particularly those that require substantial energetic investment.

2. THE PREDICTION

Many sexually selected traits depend, for their expression, on the condition of the signaller [3,10,11]. Such signals can be reliable indicators of the signaller's genetic quality because 'condition' can be influenced by a whole range of genes segregating throughout the genome [11].

* Author for correspondence (michael.garratt@unsw.edu.au).

Individuals able to acquire good condition, therefore, are more likely than average individuals to have high breeding values for fitness, including an ability to avoid infection by prevailing parasite and pathogen strains.

Von Schantz *et al.* [6] suggested that the links between oxidative stress and organism health, long recognized by physiologists, might provide one important mechanism linking condition to sexual signalling [6]. Oxidative stress is a negative, often pathological process that occurs when there is an overproduction of reactive oxygen species (ROS) in relation to an organism's ability to control their damaging effects ([8]; figure 1). ROS are either free radicals (molecules with one or more unpaired electrons, making them very reactive) or molecules with a high propensity to react with other molecules to produce free radicals. ROS can cause damage to biomolecules (termed oxidative damage), impair cellular function and even cause death [12]. As a consequence, animals have a variety of defence mechanisms that under normal physiological conditions protect against the potentially damaging effects of ROS. These include the superoxide dismutase, catalase and glutathione peroxidase families of antioxidant enzymes, and non-enzymatic antioxidants such as glutathione and dietary derived vitamin E.

We stress that there are many other antioxidant defences and protective mechanisms, and a variety of repair mechanisms that help to protect against unwanted oxidation of cellular components and maintain redox homeostasis. However, under certain conditions, these defences are ineffective and oxidative stress can result [8]. Normal cellular metabolism is expected to produce the majority of ROS in biological systems [13], as ROS can be leaked by the electron transport chains during oxidative phosphorylation. However, the immune and detoxification systems also produce ROS when activated, such that individuals with substantial activation of these systems can suffer elevated oxidative stress [6]. Oxidative stress is further correlated with the onset and/or severity of hundreds of diseases [12], increasing its applicability as a marker of condition (figure 1). Because of these links, von Schantz *et al.* [6] predicted that female choice may have selected for traits in males that reliably reveal low oxidative stress, as this would be a good marker of male quality. How and why might oxidative stress be honestly revealed by sexual signals? Von Schantz *et al.* [6] suggested a number of signals and mechanisms that could reveal this aspect of physiology, a list that other authors have added to in the intervening 13 years. While keeping in mind a modern understanding of the physiology of oxidative stress, we next very briefly summarize and evaluate those previously suggested mechanisms that have gained the most empirical attention.

(a) *Oxidative stress and visual sexual signals*

The majority of empirical and theoretical research linking oxidative stress to sexual signals has focused on species where males produce colourful carotenoid-based traits that are attractive to females [14–17]. A considerable body of research rests on the idea that because carotenoids cannot be synthesized *de novo* by animals (but see Kemp *et al.* [18]), and must be obtained solely from the diet, males with the largest and most colourful signals may be advertising their foraging ability [19,20]. Likewise, if

carotenoids also act as immune-enhancers and antioxidants [6,21], then males that can afford to tie carotenoid molecules up in visual signals might be demonstrating their good health, as they do not need to allocate these molecules to health functions. There is evidence that carotenoids are immune-enhancers and females that select males on the basis of carotenoid-based signals gain immune-competent mates [22,23], but there is much less evidence that carotenoids have an antioxidant role *in vivo* [24,25], which has caused this hypothesis to fall out of favour with evolutionary biologists.

It must also be noted that carotenoids might signal oxidative stress through other pathways. They can be oxidized and lose their coloration, so males with high oxidative stress may be unable to maintain these pigment molecules in their colourful form [26]. Manipulations of other dietary antioxidants have been shown to influence either the size [27] or intensity of [28,29] coloration of several different carotenoid-based visual signals, supporting this theory [27–29]. However, manipulation of ROS levels through treatment with the molecules paraquat or diaquat, which induce an *in vivo* ROS attack, have produced inconsistent results, with increased ROS production reducing the coloration of carotenoid-based traits in some studies [30] but not others [31]. Although carotenoids have an important role in signalling other information, their ability to signal oxidative stress seems far from ubiquitous.

More recently, the effects of oxidative stress on sexual signalling have focused on changes in melanin-based visual traits. Traditionally, melanin-based signals were expected to be under genetic control and carotenoid signals were expected to reveal condition, although there is little evidence that these signals convey different information [32]. A variety of mechanisms have been proposed that would link the expression of melanin-based traits to oxidative stress, although, as recently highlighted, these pathways and their evolution require further understanding [9]. A full discussion of these is beyond the scope of this article (see [9] and references within), but one hypothesis suggests that individuals are forced to increase their susceptibility to oxidative stress when investing in melanin-based signals [33], possibly because glutathione, an important antioxidant, must be reduced to allow maximal melanization. This handicap could ensure that only males with adequate antioxidant protection can produce these signals while coping with such a cost.

Inhibition of glutathione production in both great tits (*Parus major*) [33] and greenfinches (*Carduelis chloris*) [34] has led to increased investment in melanin-based visual sexual signals, illustrating that a reduction in glutathione levels can increase melanization. Furthermore, treatment of developing red-legged partridges (*Alectoris rufa*) with the pro-oxidant molecule diaquat also led to an increase in the size of melanin trait [35]. Current empirical research thus suggests that investment in some melanin-based traits is related to an individual's level of oxidative stress; somewhat paradoxically, however, these studies show that elevation of oxidative stress leads to an *increase* in signalling effort. It has been noted that experimental reduction of glutathione levels leads to an increase in the total antioxidant capacity (TAC) of blood [33], supporting the theory of the authors that 'melanin-based traits are in fact signals of the individual




Figure 1. How oxidative stress, an overproduction of ROS that cannot be managed by protective mechanisms, can reveal mate quality. ROS are produced from a number of different sources and their production has the potential to cause oxidative stress. However, animals have a variety of different protective mechanisms that protect against such damaging effects and often work in concert with ROS production to maintain homeostasis. Under some conditions, however, there is a breakdown in homeostasis and oxidative stress results. This can correlate with negative physiological states that in many circumstances would be undesirable in a mate. Arrows indicate causal directions, with the large arrow indicating the key factor influencing oxidative stress.

capacity to manage oxidative stress' [33]. A better understanding of the way individuals differ in their management of oxidative stress would help to reveal the precise mechanisms that maintain honesty in such systems, which is a principle aim of this article.

(b) *Testosterone-dependent and energetically costly sexual signals*

Another hypothesis that suggests individuals differ in their ability to manage oxidative stress while signalling relates to the dependence of many secondary sexual traits on elevated testosterone levels. If the required increases in testosterone have negative effects such as immune suppression, then only individuals in good health and with few parasites will be able to cope with these effects while investing in signalling [36]. The evidence for an immunosuppressive effect of testosterone is conflicting [37]. Alonso-Alvarez *et al.* [38] suggested that high levels of testosterone may instead generate oxidative stress and this may be the physiological constraint that maintains honesty. Under this scenario, oxidative stress may maintain the honesty of a range of testosterone-dependent traits, not just visual signals, although the focus of empirical work has been largely on birds and fishes.

Experimental elevation of testosterone levels decreased antioxidant capacity [38] and increased oxidative damage

[39] in zebra finches (*Taeniopygia guttata*) and red grouse (*Lagopus lagopus scoticus*) respectively. In red-legged partridges, however, elevation of testosterone did not produce oxidative damage but instead caused a decrease in the coloration of a carotenoid-based signal, highlighting that testosterone levels and sexual signalling may not always be correlated. Similarly, in house mice (*Mus musculus domesticus*), where investment in particular olfactory signals is influenced by castration and subsequent testosterone treatment [40,41], males were able to greatly increase investment in olfactory signalling over a 16-week period, while at the same time reducing both testosterone levels and oxidative damage in liver and serum [42].

Testosterone may inadvertently cause oxidative stress because it stimulates an increase in metabolic rate, which, potentially, could cause greater ROS production [38]. The assumed links between metabolic rate and ROS production require further validation, as the relationship between these is not 1:1 and sometimes large increases in metabolism can cause ROS production to decrease [43; discussed later in this article]. Nonetheless, under the assumption that metabolic rate and ROS production are positively correlated in many situations, Metcalfe & Alonso-Alvarez [9] broadened the applicability of oxidative stress in sexual signalling by suggesting that secondary sexual traits 'whose production

requires elevated oxygen consumption should be good candidates to act as signallers of good antioxidant defences'—thus even signals that are not testosterone-dependent may indicate an individual's level of oxidative stress or antioxidant quality. Both of these hypotheses appear based on the premise that an increase in metabolic rate will shift the balance between ROS production and antioxidants.

Von Shantz *et al.* saw oxidative stress as the outcome of a balance between ROS production and antioxidant defences, and little emphasis was placed on the dynamic regulation of antioxidant defences that occurs in response to a variety of stressors. The ability of organisms to regulate defence mechanisms is beginning to be appreciated by behavioural ecologists [8,44,45], but the precise mechanisms that allow such regulation are rarely discussed and almost never tested. In many empirical studies, TAC is measured as a marker of defence against oxidative stress, which essentially involves assessing the ability of a sample to quench an experimentally generated oxidant [46]. Although these assays are relatively easy to run and require small amounts of sample, they give little indication of the exact mechanisms that individuals use to control ROS production and allow future protection against oxidative stress. It is also a matter of uncertainty as to what these TAC assays actually detect. Such *in vitro* assays are conducted outside of their usual biological context, which is characterized by a greater enzymatic maintenance of steady state [46]. Furthermore, some measures of TAC preferentially detect the concentration of uric acid over other, arguably more important lines of antioxidant defence [47].

In §3, we highlight how endogenous antioxidant defences are regulated in response to transient increases in ROS production. Moreover, and central to this study, we emphasize that not all individuals have a similar capacity to regulate defences against oxidative stress and this may lead to condition-dependent costs of sexual signalling.

3. OXIDATIVE STRESS IN THE TWENTY-FIRST CENTURY

Understanding of the physiology of oxidative stress has changed substantially in the 13 years since the von Shantz *et al.* review paper. In 1999, oxidative stress was still sometimes considered to be the outcome of a 'seesaw effect' between the balance of unwanted ROS production and the quantity of antioxidants an animal has in its system. An increased consumption of dietary antioxidants was hypothesized to decrease oxidative stress and protect against disease and this has been an important component of hypotheses linking oxidative stress to the expression of colourful visual signals. However, the evidence that dietary molecules, particularly carotenoids, function as antioxidants *in vivo* is slim [24,25]. Furthermore, a number of studies in humans have suggested that an increased consumption of certain antioxidants, such as vitamins A, E and β -carotene, can actually be detrimental to health [48,49]. One possible reason for these detrimental effects is the inhibition of physiological regulatory processes that are mediated by ROS [50].

While ROS can be damaging at high concentrations, at lower levels they have wide-ranging adaptive roles and are

produced by a number of enzymes such as NADPH oxidase and nitric oxide synthase [12]. ROS are essential in the adaptive immune response and have a plethora of roles in cellular signalling. These include the control of vascular tone and platelet adhesion, oxygen sensing, cell adhesion and division, contraction and apoptosis [51,52]. Redox signalling, a regulatory process in which signals are delivered through redox chemistry, also involves ROS as essential mediators and plays an important role in protection against oxidative stress [53].

Transient increases in ROS levels can elicit the upregulation of a number of defences that protect against oxidative stress. Increased ROS production from the electron transport chains during energy metabolism can increase the expression and concentrations of specialized uncoupling proteins, which increase proton conductance across the inner mitochondrial membrane in the presence of specific activators [54]. This mild uncoupling of the mitochondria produces heat, which is an active function of one particular uncoupling protein, UCP1. However, uncoupling also reduces the leakage of electrons that give rise to ROS [55] and may be used as an adaptive mechanism to protect against oxidative stress. This may be a principle role of two other uncoupling proteins: UCP2 and UCP3 [54].

Increased ROS production can also cause an upregulation of antioxidant defences and cellular repair mechanisms that protect against oxidative stress. These changes are usually dynamic, with increased ROS causing a shift to a slightly oxidized cellular state. These changes in the redox state influence signalling pathways and gene expression, which help to increase the activities of antioxidant defences and restore redox homeostasis [52]. We give examples of changes to specific antioxidants in the following paragraph; however, it is important to note that there are a variety of discrete pathways that can be modulated to help protect against increased ROS, with upregulation of DNA and protein repair mechanisms also occurring to reduce any damage that has occurred [56]. Upregulation of protective mechanisms may have long-term beneficial effects and it has even been suggested that the known lifespan extending effect of calorie restriction may be a consequence of an increase in mitochondrial metabolism [57]. It has been hypothesized that this increase in metabolism could increase ROS production, which in turn elevates stress resistance and antioxidant defence, facilitating a longer lifespan [58]. This theory, termed mitohormesis, is at odds with the classical, outdated view of ROS as purely negative, damaging molecules.

Some of the best examples of the dynamic regulation of specific antioxidant defences derive from studies of exercise physiology in mice and humans. In mice, a single session of exercise physiologically induced muscle contraction can lead to greater production of ROS and oxidative damage [59,60]. However, healthy individuals respond to such increases by upregulating defences against oxidative stress, such as antioxidant enzymes and heat shock proteins, which protect against ROS production during future exercise bouts [61]. Similar results are seen in humans after moderate exercise [62,63]. Such changes are thought to occur, at least in part, in response to activation of redox-sensitive transcription factors such as activator protein-1 (AP-1) and nuclear factor kappa-B (NF- κ B). Consuming dietary antioxidants, previously assumed to facilitate protection against oxidative stress, can actually inhibit the activation of redox-sensitive

pathways and limit the adaptive responses to ROS production that occur during exercise [64].

Dysfunctional redox regulation is also a feature of ageing [52] and is present in individuals that are suffering persistent oxidative stress [65]. For example, at rest, the muscles of old mice show chronic activation of transcription factors AP-1 and NF- κ B. However, after a demanding bout of forced muscle contraction, there is no change in the activation of these transcription factors and also no change in antioxidant expression, which may limit the ability of old males to protect against further oxidative stress [61]. This lack of redox regulation is expected to be an important cause of the age-related loss of muscle force generation [66].

The increasing recognition of redox dysfunction, particularly in relation to pathology, has led to the suggestion that oxidative stress should be redefined as 'a disruption in redox signalling and control' [67]. Under this hypothesis, the definition of oxidative stress as an imbalance between pro-oxidants and antioxidants is ineffective because of the existence of multiple discrete redox pathways that function outside of a single equilibrium. A shift towards an oxidized redox state is associated with old age, unhealthy lifestyles and many types of diseases and malignancies [53,65]. It has even been suggested that dysfunctional redox regulation could have a greater impact on senescence than the ROS-inflicted damage that occurs to cellular components [52]. Given both the variation that is exhibited in individual redox responses and its apparent ability to influence investment in energetically demanding traits, redox regulation may have a range of roles in the signalling of male quality.

(a) Condition-dependent sexual signalling and impaired redox regulation

Dynamic regulation of antioxidant defences occurs with alteration of redox state, particularly when transient increases in ROS occur [52], and this serves to restore redox balance. This regulation is also likely to result when animals invest in demanding reproductive activities, such as investment in sexual signalling. Female mice increase the concentration of glutathione, an important antioxidant, in the liver when investing in reproduction and territory defence, and this is associated with a reduction in markers of oxidative stress in comparison with non-reproductives [68]. After testosterone levels were increased experimentally in male red-legged partridges, serum levels of glutathione and carotenoids both increased without any observed changes in oxidative damage [69]. Thus, while von Schantz *et al.* [6] predicted that increased ROS production during signalling could shift the overall balance of ROS production and antioxidant defences to an oxidizing state, current understanding suggests that this balance will simply be restored by an appropriate response from the antioxidant defence system.

Although healthy individuals modulate defences in response to oxidative insults, redox regulation in individuals that are unhealthy or in poor condition has frequently been shown to be dysfunctional, limiting the ability to adapt and protect against oxidative stress [52,65,70,71]. This could have important consequences for the costs that are incurred when individuals invest in sexual signals that shift redox balance to an oxidized state. Individuals

in good condition are likely to respond to any transient oxidation incurred from signalling with an upregulation of defence mechanisms that protect against oxidative stress, limiting the viability costs of signalling. However, individuals in poor condition could suffer sustained oxidative stress resulting in greater costs despite an equivalent signalling investment. This falls in line with Zahavi's handicap hypothesis, which predicts that high-quality individuals will pay lower marginal signalling costs when compared with those incurred by poor-quality individuals that make an equivalent investment [3,72,73]. Dysfunctional redox regulation in poor condition individuals may be an important handicap influencing the honesty of condition-dependant signals.

(b) Oxidative stress and the fitness benefits of sexual signalling

There is, however, more to honest signalling than condition-dependent handicaps. Even though differences in the costliness of sexual signals to signallers of differing quality have gathered most theoretic and empirical attention [73], Getty [74] has argued that honesty can also arise in a system where high-quality signallers benefit more from a given level of signalling than low-quality individuals. Getty [74] made this point in the context of parasite-induced costs and showed that honest signalling systems could result in either a positive or a negative relationship between individual signal expression and parasite burden. Oxidative stress and sexual signalling are usually interpreted in the context of the handicap hypothesis, which predicts a negative correlation between the expression of a signal and the handicap it imposes [9,72]. However, there are many possible honest biological signalling systems where this condition will not hold true.

Honesty in a signalling system occurs when it is in the interest of poor-quality individuals to signal at a lower level than high-quality males. Condition-dependent costs can cause signalling strategies to diverge between males that differ in quality, but differences in the fecundity benefits of signalling are equally important [1,75]. If lower quality males get a smaller fecundity return from signalling than higher quality males, it will be in their strategic interest to invest less in signalling (because the benefits are fewer) when other things are equal.

Oxidative stress can have potentially severe negative impacts on male sperm quality [7,76,77], which may decrease the fecundity benefits of signalling because males are less likely to convert any attraction of females into successful fertilizations. For example, both the percentage of motile sperm and sperm swimming ability can correlate negatively with oxidative stress [78,79], suggesting that oxidative stress may influence the functional characteristics of sperm. Various oxidative insults incurred along the female genital tract could also mean that only individual sperm free from oxidative stress and possessing good antioxidant protection will be able to reach the ovum [77]. Additionally, both these factors could cause males with high oxidative stress to be less likely to successfully fertilize a female's eggs under sperm competition [7,80]. If oxidative stress lowers the probability of a mating resulting in successful fertilization, then males with high oxidative stress may strategically

invest less in sexual signalling, as they get fewer benefits from investment in this life-history trait.

(c) *Oxidative stress and mitochondrial bioenergetics*

Many suggestions that link sexual signalling to oxidative stress are based on the assumption that increasing metabolic rate will increase ROS production. There is now evidence that the relationship between metabolic rate and ROS production is neither linear nor simple and that increases in metabolism can, in some circumstances, even reduce ROS production. As we describe later, large increases in metabolism can cause two substrates that are required for metabolic generation of ROS—electrons and oxygen—to become depleted, limiting ROS production [43].

The biggest metabolic transition that occurs with regards to ROS production is the change from state 4 to state 3 respiration. When mitochondria are in resting state 4 and ADP is absent, electrons flow slowly and tend to accumulate in the respiratory chain, causing particular complexes to be more reduced and increasing their capacity to leak electrons that subsequently produce ROS. In active state 3 respiration, when mitochondria are producing energy, the rate of electron flow strongly increases and decreases reduction of the electron transport chain, greatly reducing the probability that electrons will be leaked and therefore decreasing ROS production. In addition, during periods of high metabolism, oxygen can be depleted from areas surrounding the mitochondria, further limiting ROS production. [43]. Thus, in many situations, the assumption that an increase in metabolic rate will increase ROS production does not hold. For example, increasing the lifelong metabolic rates of short-tailed field voles (*Microtus agrestis*) by housing subjects in the cold had no effect on oxidative damage and negligible effects on antioxidant defences [81]. Furthermore, in a separate study across a population of mice, those individuals with the highest metabolic rates lived longer and had the greatest uncoupling of their muscle mitochondria [82], which is likely to have reduced ROS production. Much of the evidence linking higher metabolism to greater ROS production is derived from comparative studies across species; but within species, when an individual increases investment in signalling, the associated changes in metabolism will not always produce more ROS.

Oxidative stress may influence the ability of mitochondria to produce energy efficiently, which could have important effects on investment in sexual signalling. Many sexual signals are energetically demanding, from the energy required to generate calls or to display to the demands of carrying large ornaments or weapons around. Animals are typically expected to be limited in their available energy, and these limits may constrain investment in signalling. Indeed, one frequently used definition of ‘condition’ is the amount of resources available for allocation to fitness-enhancing traits [5]. Any changes in mitochondrial efficiency may thus have important effects on the energy available for signalling.

Recent technological advances have allowed measurement of mitochondrial energetics in intact cells [83]. Using these techniques, it has been demonstrated that oxidative stress can influence the basal respiration of mitochondria. Furthermore, mitochondria in cells under

oxidative stress have a lower ‘reserve capacity’, that is, a lower ability to increase energy production in response to cellular demands [84,85]. Changes in mitochondrial energetics in response to oxidative stress may occur because damage to cellular biomolecules depletes available cytoplasmic ATP [83]. Additionally, mitochondrial damage may limit the ability of the electron transport chain to support electron transport [83]. While the relevance of these results *in vivo* requires further validation, if oxidative stress does impact greatly on energy production, then it could severely limit investment in sexual signalling. Direct manipulations of antioxidant expression *in vivo* demonstrate that oxidative stress can effect mitochondrial energy production and subsequent investment in demanding traits that require ATP. For example, mice with a reduced expression of mitochondrial superoxide dismutase, either globally or in particular muscle fibres, show various impairments of their mitochondria that limit efficient energy production [86–90]. Furthermore, these animals have a lower endurance capacity in energetically demanding activities such as treadmill running [89,91]. Differences in blood lactate and glucose levels between knockout and wild-type mice suggest that dysfunctional mitochondrial metabolism may cause an earlier shift from aerobic to anaerobic respiration, owing to an inability to maintain ATP demand for exercise [89].

(d) *Mitochondrial dysfunction and sexual signalling*

Given the high energy costs of many sexual signals, surprisingly few studies have investigated the consequences of individual variation in mitochondrial bioenergetics for sexual signalling. Acoustic communication is heavily dependent on the energy generated from metabolism [92] and may be particularly influenced by suboptimal mitochondrial function. A recent comparative study of the metabolic costs of calling revealed that the energy cost of calling in ectotherms averages eight times the resting metabolic rate [93]. In birds, metabolic rate while calling approximately doubled, which is an equivalent increase to that of producing eggs [93].

In mammals, a variety of behaviours involved in competition for mates, including protecting females and scramble competition, have been shown to have substantial energetic costs, sometimes greater than the costs of lactation for females [94]. Andersson [10] in particular highlights that male display behaviour, such as that arising on a lek, may be limited by energy constraints, and that these constraints appear to vary between individuals. Individual differences in the ability to metabolize energy efficiently, possibly arising in part from differences in oxidative stress, could be an important determinant of an individual’s ability to invest in energetically costly sexual signals. Of course in some situations, energetic investment will be limited by an animal’s food availability; however, it has recently been highlighted that breeding animals are often not limited by food supply [95]. We suggest that empirical studies incorporating mitochondrial energetics may reveal an important physiological constraint on sexual signalling.

4. CONCLUSIONS

Over the past 13 years, understanding of oxidative stress has changed fundamentally and this has important

implications for predictions implicating oxidative stress in life-history evolution. To date, the vast majority of empirical research in this area has focused on species that signal with visual traits. Here, we highlight that this research field has the potential to be much broader, as oxidative stress could constrain investment in a variety of sexual signals of different sensory modalities.

In order to progress, we believe researchers should attend to the physiological changes that occur with investment in sexual signalling and how these differ between individuals. Instead of measuring TAC, in some circumstances, it would be much more informative to determine changes in the activities of specific antioxidants (for example the superoxide dismutases, catalases and glutathione peroxidases, to name but a few) and repair mechanisms (for example heat shock proteins, DNA excision-repair enzymes). We predict that individuals in poor condition will be less able to regulate some of these defences and protect against oxidative stress when investing in sexual signalling. Furthermore, examination of transcription factors and gene expression may give a better insight into the regulatory responses that occur with sexual signalling and whether these are impaired in individuals of poor condition. Similarly, measurements of mitochondrial energetics, either in isolated mitochondria or ideally mitochondria *in situ* (for example in permeabilized tissues, where mitochondrial function may be more similar to *in vivo* conditions; [96]), may help to reveal whether animals that differ in condition and sexual signalling also differ in their ability to produce energy efficiency. Changes in mitochondrial density could also be expected with sexual signalling, which may have important effects on both energy production and the generation of ROS; this can be measured directly (for example, by electron microscopy) and indirectly by the examination of the mitochondrial enzymes citrate synthase and cytochrome *c* oxidase [97]. Care should be taken when selecting methods to measure factors related to oxidative stress, and we recommend collaborations with physiologists to ensure accurate measurement of the trait of interest.

We have highlighted several novel pathways that could link oxidative stress to investment in a range of sexual signals associated with high-energy demands. Impaired redox regulation and mitochondrial energetics have been well studied with regard to exercise physiology and are important factors limiting endurance and muscle strength. Acoustic calling is energetically costly and investment in these signals is somewhat constrained by the size of the sound producing muscle [93]; thus, our examples from exercise physiology may be especially relevant. In mammals, investment in olfactory signalling, aggressive behaviours and displaying can all entail energetic costs and could also be influenced by oxidative stress. Only after empirical tests in more diverse animal groups, will we understand whether oxidative stress poses a general constraint on sexual signalling, as hypothesized [6,9], or whether it applies to a limited number of visual signals.

Thanks to Michael Kasumovic, Jean-François Lemaitre and Nicolas Pichaud for valuable comments on a version of the manuscript. M.G. thanks members of the Pathophysiology Research Group and Mammalian Behaviour and Evolution Group at the University of Liverpool, UK, for many interesting discussions, which

increased his understanding of this area and in doing so helped him to formulate some of the ideas presented here.

REFERENCES

- 1 Getty, T. 2006 Sexually selected signals are not similar to sports handicaps. *Trends Ecol. Evol.* **21**, 83–88. (doi:10.1016/j.tree.2005.10.016)
- 2 Maynard Smith, J. & Harper, D. 2003 *Animal signals*. Oxford, UK: Oxford University Press.
- 3 Cotton, S., Fowler, K. & Pomiankowski, A. 2004 Do sexual ornaments demonstrate heightened condition-dependent expression as predicted by the handicap hypothesis? *Proc. R. Soc. Lond. B* **271**, 771–783. (doi:10.1098/rspb.2004.2688)
- 4 Lailvaux, S. P. & Irschick, D. J. 2006 A functional perspective on sexual selection: insights and future prospects. *Anim. Behav.* **72**, 263–273. (doi:10.1016/j.anbehav.2006.02.003)
- 5 Tomkins, J. L., Radwan, J., Kotiaho, J. S. & Tregenza, T. 2004 Genic capture and resolving the lek paradox. *Trends Ecol. Evol.* **19**, 323–328. (doi:10.1016/j.tree.2004.03.029)
- 6 von Schantz, T., Bensch, S., Grahm, M., Hasselquist, D. & Wittzell, H. 1999 Good genes, oxidative stress and condition-dependent sexual signals. *Proc. R. Soc. Lond. B* **266**, 1–12. (doi:10.1098/rspb.1999.0597)
- 7 Dowling, D. K. & Simmons, L. W. 2009 Reactive oxygen species as universal constraints in life-history evolution. *Proc. R. Soc. B* **276**, 1737–1745. (doi:10.1098/rspb.2008.1791)
- 8 Monaghan, P., Metcalfe, N. B. & Torres, R. 2009 Oxidative stress as a mediator of life history trade-offs: mechanisms, measurements and interpretation. *Ecol. Lett.* **12**, 75–92. (doi:10.1111/j.1461-0248.2008.01258.x)
- 9 Metcalfe, N. B. & Alonso-Alvarez, C. 2010 Oxidative stress as a life-history constraint: the role of reactive oxygen species in shaping phenotypes from conception to death. *Funct. Ecol.* **24**, 984–996. (doi:10.1111/j.1365-2435.2010.01750.x)
- 10 Andersson, M. 1994 *Sexual selection*. New York, NY: Princeton University Press.
- 11 Rowe, L. & Houle, D. 1996 The lek paradox and the capture of genetic variance by condition dependent traits. *Proc. R. Soc. Lond. B* **263**, 1415–1421. (doi:10.1098/rspb.1996.0207)
- 12 Halliwell, B. & Gutteridge, J. M. 1999 *Free radicals in biology and medicine*. Oxford, UK: Oxford University Press.
- 13 Balaban, R. S., Nemoto, S. & Finkel, T. 2005 Mitochondria, oxidants, and aging. *Cell* **120**, 483–495. (doi:10.1016/j.cell.2005.02.001)
- 14 Olson, V. A. & Owens, I. P. F. 1998 Costly sexual signals: are carotenoids rare, risky or required? *Trends Ecol. Evol.* **13**, 510–514. (doi:10.1016/s0169-5347(98)01484-0)
- 15 Zuk, M., Thornhill, R., Ligon, J. D., Johnson, K., Austad, S., Ligon, S. H., Thornhill, N. W. & Costin, C. 1990 The role of male ornaments and courtship behavior in female mate choice of red jungle fowl. *Am. Nat.* **136**, 459–473. (doi:10.1086/285107)
- 16 Milinski, M. & Bakker, T. C. M. 1990 Female sticklebacks use male coloration in mate choice and hence avoid parasitized males. *Nature* **344**, 330–333. (doi:10.1038/344330a0)
- 17 Burley, N. & Cooper-smith, C. B. 1987 Bill color preferences of zebra finches. *Ethology* **76**, 133–151. (doi:10.1111/j.1439-0310.1987.tb00679.x)
- 18 Kemp, D. J., Herberstein, M. E. & Grether, G. F. 2012 Unraveling the true complexity of costly color signaling. *Behav. Ecol.* **23**, 233–236. (doi:10.1093/beheco/arr153)

- 19 Endler, J. A. 1980 Natural selection on color patterns in *Poecilia reticulata*. *Evolution* **34**, 76–91. (doi:10.2307/2408316)
- 20 Hill, G. E. 1992 Proximate basis of variation in carotenoid pigmentation in male house finches. *Auk* **109**, U1–U12.
- 21 Lozano, G. A. 1994 Carotenoids, parasites, and sexual selection. *Oikos* **70**, 309–311. (doi:10.2307/3545643)
- 22 Faivre, B., Gregoire, A., Preault, M., Cezilly, F. & Sorci, G. 2003 Immune activation rapidly mirrored in a secondary sexual trait. *Science* **300**, 103. (doi:10.1126/science.1081802)
- 23 Blount, J. D., Metcalfe, N. B., Birkhead, T. R. & Surai, P. F. 2003 Carotenoid modulation of immune function and sexual attractiveness in zebra finches. *Science* **300**, 125–127. (doi:10.1126/science.1082142)
- 24 Costantini, D. & Moller, A. P. 2008 Carotenoids are minor antioxidants for birds. *Funct. Ecol.* **22**, 367–370. (doi:10.1111/j.1365-2435.2007.01366.x)
- 25 Perez-Rodriguez, L. 2009 Carotenoids in evolutionary ecology: re-evaluating the antioxidant role. *Bioessays* **31**, 1116–1126. (doi:10.1002/bies.200900070)
- 26 Hartley, R. C. & Kennedy, M. W. 2004 Are carotenoids a red herring in sexual display? *Trends Ecol. Evol.* **19**, 353–354. (doi:10.1016/j.tree.2004.04.002)
- 27 Perez, C., Lores, M. & Velando, A. 2008 Availability of nonpigmentary antioxidant affects red coloration in gulls. *Behav. Ecol.* **19**, 967–973. (doi:10.1093/beheco/arn053)
- 28 Pike, T. W., Blount, J. D., Lindstrom, J. & Metcalfe, N. B. 2007 Availability of non-carotenoid antioxidants affects the expression of a carotenoid-based sexual ornament. *Biol. Lett.* **3**, 353–356. (doi:10.1098/rsbl.2007.0072)
- 29 Bertrand, S., Faivre, B. & Sorci, G. 2006 Do carotenoid-based sexual traits signal the availability of non-pigmentary antioxidants? *J. Exp. Biol.* **209**, 4414–4419. (doi:10.1242/jeb.02540)
- 30 Alonso-Alvarez, C. & Galvan, I. 2011 Free radical exposure creates paler carotenoid-based ornaments: a possible interaction in the expression of black and red traits. *PLoS ONE* **6**, e19403. (doi:10.1371/journal.pone.0019403)
- 31 Isaksson, C. & Andersson, S. 2008 Oxidative stress does not influence carotenoid mobilization and plumage pigmentation. *Proc. R. Soc. B* **275**, 309–314. (doi:10.1098/rspb.2007.1474)
- 32 Griffith, S. C., Parker, T. H. & Olson, V. A. 2006 Melanin-versus carotenoid-based sexual signals: is the difference really so black and red? *Anim. Behav.* **71**, 749–763. (doi:10.1016/j.anbehav.2005.07.016)
- 33 Galvan, I. & Alonso-Alvarez, C. 2008 An intracellular antioxidant determines the expression of a melanin-based signal in a bird. *PLoS ONE* **3**, e3335. (doi:10.1371/journal.pone.0003335)
- 34 Horak, P., Sild, E., Soomets, U., Sepp, T. & Kilk, K. 2010 Oxidative stress and information content of black and yellow plumage coloration: an experiment with greenfinches. *J. Exp. Biol.* **213**, 2225–2233. (doi:10.1242/jeb.042085)
- 35 Galvan, I. & Alonso-Alvarez, C. 2009 The expression of melanin-based plumage is separately modulated by exogenous oxidative stress and a melanocortin. *Proc. R. Soc. B* **276**, 3089–3097. (doi:10.1098/rspb.2009.0774)
- 36 Folstad, I. & Karter, A. J. 1992 Parasites, bright males, and the immunocompetence handicap. *Am. Nat.* **139**, 603–622. (doi:10.1086/285346)
- 37 Roberts, M. L., Buchanan, K. L. & Evans, M. R. 2004 Testing the immunocompetence handicap hypothesis: a review of the evidence. *Anim. Behav.* **68**, 227–239. (doi:10.1016/j.anbehav.2004.05.001)
- 38 Alonso-Alvarez, C., Bertrand, S., Faivre, B., Chastel, O. & Sorci, G. 2007 Testosterone and oxidative stress: the oxidation handicap hypothesis. *Proc. R. Soc. B* **274**, 819–825. (doi:10.1098/rspb.2006.3764)
- 39 Mougeot, F., Martinez-Padilla, J., Webster, L. M. I., Blount, J. D., Perez-Rodriguez, L. & Pieltney, S. B. 2009 Honest sexual signalling mediated by parasite and testosterone effects on oxidative balance. *Proc. R. Soc. B* **276**, 1093–1100. (doi:10.1098/rspb.2008.1570)
- 40 Kimura, T. & Hagiwara, Y. 1985 Regulation of urine marking in male and female mice: effects of sex steroids. *Horm. Behav.* **19**, 64–70. (doi:10.1016/0018-506X(85)90006-6)
- 41 Clissold, P. M., Hainey, S. & Bishop, J. O. 1984 Messenger-RNAs coding for mouse major urinary proteins are differentially induced by testosterone. *Biochem. Genet.* **22**, 379–387. (doi:10.1007/BF00484236)
- 42 Garratt, M., McArdle, F., Stockley, P., Vasilaki, A., Beynon, R. J., Jackson, M. J. & Hurst, J. L. 2011 Tissue-dependent changes in oxidative damage with male reproductive effort in house mice. *Funct. Ecol.* **26**, 423–433. (doi:10.1111/j.1365-2435.2011.01952.x)
- 43 Barja, G. 2007 Mitochondrial oxygen consumption and reactive oxygen species production are independently modulated: implications for aging studies. *Rejuvenation Res.* **10**, 215–223. (doi:10.1089/rej.2006.0516)
- 44 Costantini, D. 2010 Redox physiology in animal function: the struggle of living in an oxidant environment. *Curr. Zool.* **56**, 687–702.
- 45 Hill, G. E. 2011 Condition-dependent traits as signals of the functionality of vital cellular processes. *Ecol. Lett.* **14**, 625–634. (doi:10.1111/j.1461-0248.2011.01622.x)
- 46 Sies, H. 2007 Total antioxidant capacity: appraisal of a concept. *J. Nutr.* **137**, 1493–1495.
- 47 Costantini, D. 2011 On the measurement of circulating antioxidant capacity and the nightmare of uric acid. *Methods Ecol. Evol.* **2**, 321–325. (doi:10.1111/j.2041-210X.2010.00080.x)
- 48 Bjelakovic, G., Nikolova, D., Simonetti, R. G. & Gluud, C. 2004 Antioxidant supplements for prevention of gastrointestinal cancers: a systematic review and meta-analysis. *Lancet* **364**, 1219–1228. (doi:10.1016/S0140-6736(04)17138-9)
- 49 Seifried, H. E., Anderson, D. E., Fisher, E. I. & Milner, J. A. 2007 A review of the interaction among dietary antioxidants and reactive oxygen species. *J. Nutr. Biochem.* **18**, 567–579. (doi:10.1016/j.jnutbio.2006.10.007)
- 50 Bjelakovic, G. & Gluud, C. 2007 Surviving antioxidant supplements. *J. Natl Cancer Inst.* **99**, 742–743. (doi:10.1093/jnci/djk211)
- 51 Janssen-Heininger, Y. M. W., Mossman, B. T., Heintz, N. H., Forman, H. J., Kalyanaraman, B., Finkel, T., Stamler, J. S., Rhee, S. G. & van der Vliet, A. 2008 Redox-based regulation of signal transduction: principles, pitfalls, and promises. *Free Radical Biol. Med.* **45**, 1–17. (doi:10.1016/j.freeradbiomed.2008.03.011)
- 52 Droge, W. 2002 Free radicals in the physiological control of cell function. *Physiol. Rev.* **82**, 47–95.
- 53 Valko, M., Leibfritz, D., Moncol, J., Cronin, M. T. D., Mazur, M. & Telser, J. 2007 Free radicals and antioxidants in normal physiological functions and human disease. *Int. J. Biochem. Cell Biol.* **39**, 44–84. (doi:10.1016/j.biocel.2006.07.001)
- 54 Brand, M. D. & Esteves, T. C. 2005 Physiological functions of the mitochondrial uncoupling proteins UCP2 and UCP3. *Cell Metab.* **2**, 85–93. (doi:10.1016/j.cmet.2005.06.002)
- 55 Brand, M. D., Affourtit, C., Esteves, T. C., Green, K., Lambert, A. J., Miwa, S., Pakay, J. L. & Parker, N. 2004 Mitochondrial superoxide: production, biological

- effects, and activation of uncoupling proteins. *Free Radical Biol. Med.* **37**, 755–767. (doi:10.1016/j.freeradbiomed.2004.05.034)
- 56 Davies, K. J. A. 2000 Oxidative stress, antioxidant defenses, and damage removal, repair, and replacement systems. *IUBMB Life* **50**, 279–289. (doi:10.1080/15216540051081010)
- 57 Nisoli, E. *et al.* 2005 Calorie restriction promotes mitochondrial biogenesis by inducing the expression of eNOS. *Science* **310**, 314–317. (doi:10.1126/science.1117728)
- 58 Ristow, M. & Zarse, K. 2010 How increased oxidative stress promotes longevity and metabolic health: the concept of mitochondrial hormesis (mitohormesis). *Exp. Gerontol.* **45**, 410–418. (doi:10.1016/j.exger.2010.03.014)
- 59 Close, G. L., Kayani, A. C., Ashton, T., McArdle, A. & Jackson, M. J. 2007 Release of superoxide from skeletal muscle of adult and old mice: an experimental test of the reductive hotspot hypothesis. *Aging Cell* **6**, 189–195. (doi:10.1111/j.1474-9726.2007.00277.x)
- 60 Vasilaki, A., Mansouri, A., Van Remmen, H., van der Meulen, J. H., Larkin, L., Richardson, A. G., McArdle, A., Faulkner, J. A. & Jackson, M. J. 2006 Free radical generation by skeletal muscle of adult and old mice: effect of contractile activity. *Aging Cell* **5**, 109–117. (doi:10.1111/j.1474-9726.2006.00198.x)
- 61 Vasilaki, A., McArdle, F., Iwanejko, L. M. & McArdle, A. 2006 Adaptive responses of mouse skeletal muscle to contractile activity: the effect of age. *Mech. Ageing Dev.* **127**, 830–839. (doi:10.1016/j.mad.2006.08.004)
- 62 Gomez-Cabrera, M. C., Domenech, E. & Vina, J. 2008 Moderate exercise is an antioxidant: upregulation of antioxidant genes by training. *Free Radical Biol. Med.* **44**, 126–131. (doi:10.1016/j.freeradbiomed.2007.02.001)
- 63 Khassaf, M., Child, R. B., McArdle, A., Brodie, D. A., Esanu, C. & Jackson, M. J. 2001 Time course of responses of human skeletal muscle to oxidative stress induced by nondamaging exercise. *J. Appl. Physiol.* **90**, 1031–1035.
- 64 Ristow, M., Zarse, K., Oberbach, A., Kloting, N., Birringer, M., Kiehnopf, M., Stumvoll, M., Kahn, C. R. & Bluher, M. 2009 Antioxidants prevent health-promoting effects of physical exercise in humans. *Proc. Natl Acad. Sci. USA* **106**, 8665–8670. (doi:10.1073/pnas.0903485106)
- 65 Vasilaki, A. *et al.* 2010 The age-related failure of adaptive responses to contractile activity in skeletal muscle is mimicked in young mice by deletion of Cu, Zn superoxide dismutase. *Aging Cell* **9**, 979–990. (doi:10.1111/j.1474-9726.2010.00635.x)
- 66 Broome, C. S., Kayani, A. C., Palomero, J., Dillmann, W. H., Mestri, R., Jackson, M. J. & McArdle, A. 2006 Effect of lifelong overexpression of HSP70 in skeletal muscle on age-related oxidative stress and adaptation after nondamaging contractile activity. *FASEB J.* **20**, 1549–1551. (doi:10.1096/fj.05-4935fje)
- 67 Jones, D. P. 2006 Redefining oxidative stress. *Antioxidants Redox Signal.* **8**, 1865–1879. (doi:10.1089/ars.2006.8.1865)
- 68 Garratt, M., Vasilaki, A., Stockley, P., McArdle, F., Jackson, M. & Hurst, J. L. 2011 Is oxidative stress a physiological cost of reproduction? An experimental test in house mice. *Proc. R. Soc. B* **278**, 1098–1106. (doi:10.1098/rspb.2010.1818)
- 69 Alonso-Alvarez, C., Perez-Rodriguez, L., Mateo, R., Chastel, O. & Vinuela, J. 2008 The oxidation handicap hypothesis and the carotenoid allocation trade-off. *J. Evol. Biol.* **21**, 1789–1797. (doi:10.1111/j.1420-9101.2008.01591.x)
- 70 Jones, D. P. & Go, Y. M. 2010 Redox compartmentalization and cellular stress. *Diabetes Obes. Metab.* **12**, 116–125. (doi:10.1111/j.1463-1326.2010.01266.x)
- 71 Heo, J. Y. 2011 Redox control of GTPases: from molecular mechanisms to functional significance in health and disease. *Antioxidants Redox Signal.* **14**, 689–724. (doi:10.1089/ars.2009.2984)
- 72 Zahavi, A. 1975 Mate selection: selection for a handicap. *J. Theor. Biol.* **53**, 205–214. (doi:10.1016/0022-5193(75)90111-3)
- 73 Grafen, A. 1990 Biological signals as handicaps. *J. Theor. Biol.* **144**, 517–546. (doi:10.1016/S0022-5193(05)80088-8)
- 74 Getty, T. 2002 Signaling health versus parasites. *Am. Nat.* **159**, 363–371. (doi:10.1086/338992)
- 75 Getty, T. 1998 Handicap signalling: when fecundity and viability do not add up. *Anim. Behav.* **56**, 127–130. (doi:10.1006/anbe.1998.0744)
- 76 Blount, J. D., Moller, A. P. & Houston, D. C. 2001 Antioxidants, showy males and sperm quality. *Ecol. Lett.* **4**, 393–396. (doi:10.1046/j.1461-0248.2001.00255.x)
- 77 Velando, A., Torres, R. & Alonso-Alvarez, C. 2008 Avoiding bad genes: oxidatively damaged DNA in germ line and mate choice. *Bioessays* **30**, 1212–1219. (doi:10.1002/bies.20838)
- 78 Jones, R., Mann, T. & Sherins, R. 1979 Peroxidative breakdown of phospholipids in human spermatozoa, spermicidal properties of fatty acid peroxides, and protective action of seminal plasma. *Fertil. Steril.* **31**, 531–537.
- 79 Helfenstein, F., Losdat, S., Moller, A. P., Blount, J. D. & Richner, H. 2010 Sperm of colourful males are better protected against oxidative stress. *Ecol. Lett.* **13**, 213–222. (doi:10.1111/j.1461-0248.2009.01419.x)
- 80 Parker, G. A. 1970 Sperm competition and its evolutionary consequences in insects. *Biol. Rev. Camb. Phil. Soc.* **45**, 525–567. (doi:10.1111/j.1469-185X.1970.tb01176.x)
- 81 Selman, C., McLaren, J. S., Collins, A. R., Duthie, G. G. & Speakman, J. R. 2008 The impact of experimentally elevated energy expenditure on oxidative stress and lifespan in the short-tailed field vole *Microtus agrestis*. *Proc. R. Soc. B* **275**, 1907–1916. (doi:10.1098/rspb.2008.0355)
- 82 Speakman, J. R. *et al.* 2004 Uncoupled and surviving: individual mice with high metabolism have greater mitochondrial uncoupling and live longer. *Aging Cell* **3**, 87–95. (doi:10.1111/j.1474-9728.2004.00097.x)
- 83 Sansbury, B. E., Jones, S. P., Riggs, D. W., Darley-Usmar, V. M. & Hill, B. G. 2011 Bioenergetic function in cardiovascular cells: the importance of the reserve capacity and its biological regulation. *Chem. Biol. Interact.* **191**, 288–295. (doi:10.1016/j.cbi.2010.12.002)
- 84 Hill, B. G., Dranka, B. P., Zou, L. Y., Chatham, J. C. & Darley-Usmar, V. M. 2009 Importance of the bioenergetic reserve capacity in response to cardiomyocyte stress induced by 4-hydroxynonenal. *Biochem. J.* **424**, 99–107. (doi:10.1042/bj20090934)
- 85 Dranka, B. P., Hill, B. G. & Darley-Usmar, V. M. 2010 Mitochondrial reserve capacity in endothelial cells: the impact of nitric oxide and reactive oxygen species. *Free Rad. Biol. Med.* **48**, 905–914. (doi:10.1016/j.freeradbiomed.2010.01.015)
- 86 Kokoszka, J. E., Coskun, P., Esposito, L. A. & Wallace, D. C. 2001 Increased mitochondrial oxidative stress in the Sod2 (+/−) mouse results in the age-related decline of mitochondrial function culminating in increased apoptosis. *Proc. Natl Acad. Sci. USA* **98**, 2278–2283. (doi:10.1073/pnas.051627098)
- 87 van Remmen, H., Williams, M. D., Guo, Z. M., Estlack, L., Yang, H., Carlson, E. J., Epstein, C. J., Huang, T. T. &

- Richardson, A. 2001 Knockout mice heterozygous for Sod2 show alterations in cardiac mitochondrial function and apoptosis. *Am. J. Physiol. Heart Circul. Physiol.* **281**, H1422–H1432.
- 88 Williams, M. D., Van Remmen, H., Conrad, C. C., Huang, T. T., Epstein, C. J. & Richardson, A. 1998 Increased oxidative damage is correlated to altered mitochondrial function in heterozygous manganese superoxide dismutase knockout mice. *J. Biol. Chem.* **273**, 28 510–28 515. (doi:10.1074/jbc.273.43.28510)
- 89 Lustgarten, M. S. *et al.* 2009 Conditional knockout of Mn-SOD targeted to type IIB skeletal muscle fibers increases oxidative stress and is sufficient to alter aerobic exercise capacity. *Am. J. Physiol. Cell Physiol.* **297**, C1520–C1532. (doi:10.1152/ajpcell.00372.2009)
- 90 Lustgarten, M. S. *et al.* 2011 MnSOD deficiency results in elevated oxidative stress and decreased mitochondrial function but does not lead to muscle atrophy during aging. *Aging Cell* **10**, 493–505. (doi:10.1111/j.1474-9726.2011.00695.x)
- 91 Kinugawa, S., Wang, Z. P., Kaminski, P. M., Wolin, M. S., Edwards, J. G., Kaley, G. & Hintze, T. H. 2005 Limited exercise capacity in heterozygous manganese superoxide dismutase gene-knockout mice: roles of superoxide anion and nitric oxide. *Circulation* **111**, 1480–1486. (doi:10.1161/01.cir.0000159261.11520.63)
- 92 Gillooly, J. F. & Ophir, A. G. 2010 The energetic basis of acoustic communication. *Proc. R. Soc. B* **277**, 1325–1331. (doi:10.1098/rspb.2009.2134)
- 93 Ophir, A. G., Schrader, S. B. & Gillooly, J. F. 2010 Energetic cost of calling: general constraints and species-specific differences. *J. Evol. Biol.* **23**, 1564–1569. (doi:10.1111/j.1420-9101.2010.02005.x)
- 94 Lane, J. E., Boutin, S., Speakman, J. R. & Humphries, M. M. 2010 Energetic costs of male reproduction in a scramble competition mating system. *J. Anim. Ecol.* **79**, 27–34. (doi:10.1111/j.1365-2656.2009.01592.x)
- 95 Speakman, J. R. & Krol, E. 2010 The heat dissipation limit theory and evolution of life histories in endotherms: time to dispose of the disposable soma theory? *Integr. Comp. Biol.* **50**, 793–807. (doi:10.1093/icb/icq049)
- 96 Picard, M., Ritchie, D., Wright, K. J., Romestaing, C., Thomas, M. M., Rowan, S. L., Taivassalo, T. & Hepple, R. T. 2010 Mitochondrial functional impairment with aging is exaggerated in isolated mitochondria compared to permeabilized myofibers. *Aging Cell* **9**, 1032–1046. (doi:10.1111/j.1474-9726.2010.00628.x)
- 97 Magwere, T., Goodall, S., Skepper, J., Mair, W., Brand, M. D. & Partridge, L. 2006 The effect of dietary restriction on mitochondrial protein density and flight muscle mitochondrial morphology in *Drosophila*. *J. Gerontol. A Biol. Sci. Med. Sci.* **61**, 36–47. (doi:10.1093/gerona/61.1.36)